

# Search for new phenomena in events with two Z bosons and missing transverse momentum in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV

T. Aaltonen,<sup>22</sup> B. Álvarez González,<sup>10,aa</sup> S. Amerio,<sup>41a</sup> D. Amidei,<sup>33</sup> A. Anastassov,<sup>16,y</sup> A. Annovi,<sup>18</sup> J. Antos,<sup>13</sup> G. Apollinari,<sup>16</sup> J. A. Appel,<sup>16</sup> T. Arisawa,<sup>55</sup> A. Artikov,<sup>14</sup> J. Asaadi,<sup>50</sup> W. Ashmanskas,<sup>16</sup> B. Auerbach,<sup>58</sup> A. Aurisano,<sup>50</sup> F. Azfar,<sup>40</sup> W. Badgett,<sup>16</sup> T. Bae,<sup>26</sup> A. Barbaro-Galtieri,<sup>27</sup> V. E. Barnes,<sup>45</sup> B. A. Barnett,<sup>24</sup> P. Barria,<sup>43a,41b</sup> P. Bartos,<sup>13</sup> M. Bause,<sup>41a,41b</sup> F. Bedeschi,<sup>43a</sup> S. Behari,<sup>24</sup> G. Bellettini,<sup>43a,43b</sup> J. Bellinger,<sup>57</sup> D. Benjamin,<sup>15</sup> A. Beretvas,<sup>16</sup> A. Bhatti,<sup>47</sup> D. Bisello,<sup>41a,41b</sup> I. Bizjak,<sup>29</sup> K. R. Bland,<sup>5</sup> B. Blumenfeld,<sup>24</sup> A. Bocci,<sup>15</sup> A. Bodek,<sup>46</sup> D. Bortoletto,<sup>45</sup> J. Boudreau,<sup>44</sup> A. Boveia,<sup>12</sup> L. Brigliadori,<sup>6a,6b</sup> C. Bromberg,<sup>34</sup> E. Brucken,<sup>22</sup> J. Budagov,<sup>14</sup> H. S. Budd,<sup>46</sup> K. Burkett,<sup>16</sup> G. Busetto,<sup>41a,41b</sup> P. Bussey,<sup>20</sup> A. Buzatu,<sup>32</sup> A. Calamba,<sup>11</sup> C. Calancha,<sup>30</sup> S. Camarda,<sup>4</sup> M. Campanelli,<sup>29</sup> M. Campbell,<sup>33</sup> F. Canelli,<sup>11,16</sup> B. Carls,<sup>23</sup> D. Carlsmith,<sup>57</sup> R. Carosi,<sup>43a</sup> S. Carrillo,<sup>17,n</sup> S. Carron,<sup>16</sup> B. Casal,<sup>10,1</sup> M. Casarsa,<sup>51a</sup> A. Castro,<sup>6a,6b</sup> P. Catastini,<sup>21</sup> D. Cauz,<sup>51a</sup> V. Cavaliere,<sup>23</sup> M. Cavalli-Sforza,<sup>4</sup> A. Cerri,<sup>27,g</sup> L. Cerrito,<sup>29,t</sup> Y. C. Chen,<sup>1</sup> M. Chertok,<sup>7</sup> G. Chiarelli,<sup>43a</sup> G. Chlachidze,<sup>16</sup> F. Chlebana,<sup>16</sup> K. Cho,<sup>26</sup> D. Chokheli,<sup>14</sup> W. H. Chung,<sup>57</sup> Y. S. Chung,<sup>46</sup> M. A. Ciocci,<sup>43a,43c</sup> A. Clark,<sup>19</sup> C. Clarke,<sup>56</sup> G. Compostella,<sup>41a,41b</sup> M. E. Convery,<sup>16</sup> J. Conway,<sup>7</sup> M. Corbo,<sup>16</sup> M. Cordelli,<sup>18</sup> C. A. Cox,<sup>7</sup> D. J. Cox,<sup>7</sup> F. Crescioli,<sup>43a,43b</sup> J. Cuevas,<sup>10,aa</sup> R. Culbertson,<sup>16</sup> D. Dagenhart,<sup>16</sup> N. d'Ascenzo,<sup>16,x</sup> M. Datta,<sup>16</sup> P. de Barbaro,<sup>46</sup> M. Dell'Orso,<sup>43a,43b</sup> L. Demortier,<sup>47</sup> M. Deninno,<sup>6a</sup> F. Devoto,<sup>22</sup> M. d'Errico,<sup>41a,41b</sup> A. Di Canto,<sup>43a,43b</sup> B. Di Ruzza,<sup>16</sup> J. R. Dittmann,<sup>5</sup> M. D'Onofrio,<sup>28</sup> S. Donati,<sup>43a,43b</sup> P. Dong,<sup>16</sup> M. Dorigo,<sup>51a</sup> T. Dorigo,<sup>41a</sup> K. Ebina,<sup>55</sup> A. Elagin,<sup>50</sup> A. Eppig,<sup>33</sup> R. Erbacher,<sup>7</sup> S. Errede,<sup>23</sup> N. Ershaidat,<sup>16,ee</sup> R. Eusebi,<sup>50</sup> S. Farrington,<sup>40</sup> M. Feindt,<sup>25</sup> J. P. Fernandez,<sup>30</sup> R. Field,<sup>17</sup> G. Flanagan,<sup>16,v</sup> R. Forrest,<sup>7</sup> M. J. Frank,<sup>5</sup> M. Franklin,<sup>21</sup> J. C. Freeman,<sup>16</sup> Y. Funakoshi,<sup>55</sup> I. Furic,<sup>17</sup> M. Gallinaro,<sup>47</sup> J. E. Garcia,<sup>19</sup> A. F. Garfinkel,<sup>45</sup> P. Garosi,<sup>43a,43c</sup> H. Gerberich,<sup>23</sup> E. Gerchtein,<sup>16</sup> S. Giagu,<sup>48a</sup> V. Giakoumopoulou,<sup>3</sup> P. Giannetti,<sup>43a</sup> K. Gibson,<sup>44</sup> C. M. Ginsburg,<sup>16</sup> N. Giokaris,<sup>3</sup> P. Giromini,<sup>18</sup> G. Giurgiu,<sup>24</sup> V. Glagolev,<sup>14</sup> D. Glenzinski,<sup>16</sup> M. Gold,<sup>36</sup> D. Goldin,<sup>50</sup> N. Goldschmidt,<sup>17</sup> A. Golossanov,<sup>16</sup> G. Gomez,<sup>10</sup> G. Gomez-Ceballos,<sup>31</sup> M. Goncharov,<sup>31</sup> O. González,<sup>30</sup> I. Gorelov,<sup>36</sup> A. T. Goshaw,<sup>15</sup> K. Goulianos,<sup>47</sup> S. Grinstein,<sup>4</sup> C. Grosso-Pilcher,<sup>12</sup> R. C. Group,<sup>53,16</sup> J. Guimaraes da Costa,<sup>21</sup> S. R. Hahn,<sup>16</sup> E. Halkiadakis,<sup>49</sup> A. Hamaguchi,<sup>39</sup> J. Y. Han,<sup>46</sup> F. Happacher,<sup>18</sup> K. Hara,<sup>52</sup> D. Hare,<sup>49</sup> M. Hare,<sup>53</sup> R. F. Harr,<sup>56</sup> K. Hatakeyama,<sup>5</sup> C. Hays,<sup>40</sup> M. Heck,<sup>25</sup> J. Heinrich,<sup>42</sup> M. Herndon,<sup>57</sup> S. Hewamanage,<sup>5</sup> A. Hocker,<sup>16</sup> W. Hopkins,<sup>16,h</sup> D. Horn,<sup>25</sup> S. Hou,<sup>1</sup> R. E. Hughes,<sup>37</sup> M. Hurwitz,<sup>12</sup> U. Husemann,<sup>58</sup> N. Hussain,<sup>32</sup> M. Hussein,<sup>34</sup> J. Huston,<sup>34</sup> G. Introzzi,<sup>43a</sup> M. Iori,<sup>48a,48b</sup> A. Ivanov,<sup>7,q</sup> E. James,<sup>16</sup> D. Jang,<sup>11</sup> B. Jayatilaka,<sup>15</sup> E. J. Jeon,<sup>26</sup> S. Jindariani,<sup>16</sup> M. Jones,<sup>45</sup> K. K. Joo,<sup>26</sup> S. Y. Jun,<sup>11</sup> T. R. Junk,<sup>16</sup> T. Kamon,<sup>25,50</sup> P. E. Karchin,<sup>56</sup> A. Kasmi,<sup>5</sup> Y. Kato,<sup>39,p</sup> W. Ketchum,<sup>12</sup> J. Keung,<sup>42</sup> V. Khotilovich,<sup>50</sup> B. Kilminster,<sup>16</sup> D. H. Kim,<sup>26</sup> H. S. Kim,<sup>26</sup> J. E. Kim,<sup>26</sup> M. J. Kim,<sup>18</sup> S. B. Kim,<sup>26</sup> S. H. Kim,<sup>52</sup> Y. K. Kim,<sup>12</sup> Y. J. Kim,<sup>26</sup> N. Kimura,<sup>55</sup> M. Kirby,<sup>16</sup> S. Klimenko,<sup>17</sup> K. Knoepfel,<sup>16</sup> K. Kondo,<sup>55,a</sup> D. J. Kong,<sup>26</sup> J. Konigsberg,<sup>17</sup> A. V. Kotwal,<sup>15</sup> M. Kreps,<sup>25</sup> J. Kroll,<sup>42</sup> D. Krop,<sup>12</sup> M. Kruse,<sup>15</sup> V. Krutelyov,<sup>50,d</sup> T. Kuhr,<sup>25</sup> M. Kurata,<sup>52</sup> S. Kwang,<sup>12</sup> A. T. Laasanen,<sup>45</sup> S. Lami,<sup>43a</sup> S. Lammel,<sup>16</sup> M. Lancaster,<sup>29</sup> R. L. Lander,<sup>7</sup> K. Lannon,<sup>37,z</sup> A. Lath,<sup>49</sup> G. Latino,<sup>43a,43c</sup> T. LeCompte,<sup>2</sup> E. Lee,<sup>50</sup> H. S. Lee,<sup>12,r</sup> J. S. Lee,<sup>26</sup> S. W. Lee,<sup>50,cc</sup> S. Leo,<sup>43a,43b</sup> S. Leone,<sup>43a</sup> J. D. Lewis,<sup>16</sup> A. Limosani,<sup>15,u</sup> C.-J. Lin,<sup>27</sup> M. Lindgren,<sup>16</sup> E. Lipeles,<sup>42</sup> A. Lister,<sup>19</sup> D. O. Litvintsev,<sup>16</sup> C. Liu,<sup>44</sup> H. Liu,<sup>54</sup> Q. Liu,<sup>45</sup> T. Liu,<sup>16</sup> S. Lockwitz,<sup>58</sup> A. Loginov,<sup>58</sup> D. Lucchesi,<sup>41a,41b</sup> J. Lueck,<sup>25</sup> P. Lujan,<sup>27</sup> P. Lukens,<sup>16</sup> G. Lungu,<sup>47</sup> J. Lys,<sup>27</sup> R. Lysak,<sup>13,f</sup> R. Madrak,<sup>16</sup> K. Maeshima,<sup>16</sup> P. Maestro,<sup>43a,43c</sup> S. Malik,<sup>47</sup> G. Manca,<sup>28,b</sup> A. Manousakis-Katsikakis,<sup>3</sup> F. Margaroli,<sup>48a</sup> C. Marino,<sup>25</sup> M. Martínez,<sup>4</sup> P. Mastrandrea,<sup>48a</sup> K. Matera,<sup>23</sup> M. E. Mattson,<sup>56</sup> A. Mazzacane,<sup>16</sup> P. Mazzanti,<sup>6a</sup> K. S. McFarland,<sup>46</sup> P. McIntyre,<sup>50</sup> R. McNulty,<sup>28,k</sup> A. Mehta,<sup>28</sup> P. Mehtala,<sup>22</sup> C. Mesropian,<sup>47</sup> T. Miao,<sup>16</sup> D. Mietlicki,<sup>33</sup> A. Mitra,<sup>1</sup> H. Miyake,<sup>52</sup> S. Moed,<sup>16</sup> N. Moggi,<sup>6a</sup> M. N. Mondragon,<sup>16,n</sup> C. S. Moon,<sup>26</sup> R. Moore,<sup>16</sup> M. J. Morello,<sup>43a,43d</sup> J. Morlock,<sup>25</sup> P. Movilla Fernandez,<sup>16</sup> A. Mukherjee,<sup>16</sup> Th. Muller,<sup>25</sup> P. Murat,<sup>16</sup> M. Mussini,<sup>6a,6b</sup> J. Nachtman,<sup>16,o</sup> Y. Nagai,<sup>52</sup> J. Naganoma,<sup>55</sup> I. Nakano,<sup>38</sup> A. Napier,<sup>53</sup> J. Nett,<sup>50</sup> C. Neu,<sup>54</sup> M. S. Neubauer,<sup>23</sup> J. Nielsen,<sup>27,e</sup> L. Nodulman,<sup>2</sup> S. Y. Noh,<sup>26</sup> O. Norniella,<sup>23</sup> L. Oakes,<sup>40</sup> S. H. Oh,<sup>15</sup> Y. D. Oh,<sup>26</sup> I. Oksuzian,<sup>54</sup> T. Okusawa,<sup>39</sup> R. Orava,<sup>22</sup> L. Ortolan,<sup>4</sup> S. Pagan Griso,<sup>41a,41b</sup> C. Pagliarone,<sup>51a</sup> E. Palencia,<sup>10,g</sup> V. Papadimitriou,<sup>16</sup> A. A. Paramonov,<sup>2</sup> J. Patrick,<sup>16</sup> G. Pauletta,<sup>51a,51b</sup> M. Paulini,<sup>11</sup> C. Paus,<sup>31</sup> D. E. Pellett,<sup>7</sup> A. Penzo,<sup>51a</sup> T. J. Phillips,<sup>15</sup> G. Piacentino,<sup>43a</sup> E. Pianori,<sup>42</sup> J. Pilot,<sup>37</sup> K. Pitts,<sup>23</sup> C. Plager,<sup>9</sup> L. Pondrom,<sup>57</sup> S. Poprocki,<sup>16,h</sup> K. Potamianos,<sup>45</sup> F. Prokoshin,<sup>14,dd</sup> A. Pranko,<sup>27</sup> F. Pthos,<sup>18,i</sup> G. Punzi,<sup>43a,43b</sup> A. Rahaman,<sup>44</sup> V. Ramakrishnan,<sup>52</sup> N. Ranjan,<sup>45</sup> I. Redondo,<sup>30</sup> P. Renton,<sup>40</sup> M. Rescigno,<sup>48a</sup> T. Riddick,<sup>29</sup> F. Rimondi,<sup>6a,6b</sup> L. Ristori,<sup>42,16</sup> A. Robson,<sup>20</sup> T. Rodrigo,<sup>10</sup> T. Rodriguez,<sup>42</sup> E. Rogers,<sup>23</sup> S. Rolli,<sup>53,j</sup> R. Roser,<sup>16</sup> F. Ruffini,<sup>43a,43c</sup> A. Ruiz,<sup>10</sup> J. Russ,<sup>11</sup> V. Rusu,<sup>16</sup> A. Safonov,<sup>50</sup> W. K. Sakumoto,<sup>46</sup> Y. Sakurai,<sup>55</sup> L. Santi,<sup>51a,51b</sup> K. Sato,<sup>52</sup> V. Saveliev,<sup>16,x</sup> A. Savoy-Navarro,<sup>16,bb</sup> P. Schlabach,<sup>16</sup> A. Schmidt,<sup>25</sup> E. E. Schmidt,<sup>16</sup> T. Schwarz,<sup>16</sup> L. Scodellaro,<sup>10</sup> A. Scribano,<sup>43a,43c</sup> F. Scuri,<sup>43a</sup> S. Seidel,<sup>36</sup> Y. Seiya,<sup>39</sup> A. Semenov,<sup>14</sup> F. Sforza,<sup>43a,43c</sup> S. Z. Shalhout,<sup>7</sup>

T. Shears,<sup>28</sup> P. F. Shepard,<sup>44</sup> M. Shimojima,<sup>52,w</sup> M. Shochet,<sup>12</sup> I. Shreyber-Tecker,<sup>35</sup> A. Simonenko,<sup>14</sup> P. Sinervo,<sup>32</sup> K. Sliwa,<sup>53</sup> J. R. Smith,<sup>7</sup> F. D. Snider,<sup>16</sup> A. Soha,<sup>16</sup> V. Sorin,<sup>4</sup> H. Song,<sup>44</sup> P. Squillacioti,<sup>43a,43c</sup> M. Stancari,<sup>16</sup> R. St. Denis,<sup>20</sup> B. Stelzer,<sup>32</sup> O. Stelzer-Chilton,<sup>32</sup> D. Stentz,<sup>16,y</sup> J. Strologas,<sup>36</sup> G. L. Strycker,<sup>33</sup> Y. Sudo,<sup>52</sup> A. Sukhanov,<sup>16</sup> I. Suslov,<sup>14</sup> K. Takemasa,<sup>52</sup> Y. Takeuchi,<sup>52</sup> J. Tang,<sup>12</sup> M. Tecchio,<sup>33</sup> P. K. Teng,<sup>1</sup> J. Thom,<sup>16,h</sup> J. Thome,<sup>11</sup> G. A. Thompson,<sup>23</sup> E. Thomson,<sup>42</sup> D. Toback,<sup>50</sup> S. Tokar,<sup>13</sup> K. Tollefson,<sup>34</sup> T. Tomura,<sup>52</sup> D. Tonelli,<sup>16</sup> S. Torre,<sup>18</sup> D. Torretta,<sup>16</sup> P. Totaro,<sup>41a</sup> M. Trovato,<sup>43a,43d</sup> F. Ukegawa,<sup>52</sup> S. Uozumi,<sup>26</sup> A. Varganov,<sup>33</sup> F. Vázquez,<sup>17,n</sup> J. Vasquez,<sup>8</sup> G. Velev,<sup>16</sup> C. Vellidis,<sup>16</sup> M. Vidal,<sup>45</sup> I. Vila,<sup>10</sup> R. Vilar,<sup>10</sup> J. Vizán,<sup>10</sup> M. Vogel,<sup>36</sup> G. Volpi,<sup>18</sup> P. Wagner,<sup>42</sup> R. L. Wagner,<sup>16</sup> T. Wakisaka,<sup>39</sup> R. Wallny,<sup>9</sup> S. M. Wang,<sup>1</sup> A. Warburton,<sup>32</sup> D. Waters,<sup>29</sup> W. C. Wester III,<sup>16</sup> D. Whiteson,<sup>42,c</sup> A. B. Wicklund,<sup>2</sup> E. Wicklund,<sup>16</sup> S. Wilbur,<sup>12</sup> F. Wick,<sup>25</sup> H. H. Williams,<sup>42</sup> J. S. Wilson,<sup>37</sup> P. Wilson,<sup>16</sup> B. L. Winer,<sup>37</sup> P. Wittich,<sup>16,h</sup> S. Wolbers,<sup>16</sup> H. Wolfe,<sup>37</sup> T. Wright,<sup>33</sup> X. Wu,<sup>19</sup> Z. Wu,<sup>5</sup> K. Yamamoto,<sup>39</sup> D. Yamato,<sup>39</sup> T. Yang,<sup>16</sup> U. K. Yang,<sup>12,s</sup> Y. C. Yang,<sup>26</sup> W.-M. Yao,<sup>27</sup> G. P. Yeh,<sup>16</sup> K. Yi,<sup>16,o</sup> J. Yoh,<sup>16</sup> K. Yorita,<sup>55</sup> T. Yoshida,<sup>39,m</sup> G. B. Yu,<sup>15</sup> I. Yu,<sup>26</sup> S. S. Yu,<sup>16</sup> J. C. Yun,<sup>16</sup> A. Zanetti,<sup>51a</sup> Y. Zeng,<sup>15</sup> C. Zhou,<sup>15</sup> and S. Zucchelli<sup>6a,6b</sup>

(CDF Collaboration)

<sup>1</sup>*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*

<sup>2</sup>*Argonne National Laboratory, Argonne, Illinois 60439, USA*

<sup>3</sup>*University of Athens, 157 71 Athens, Greece*

<sup>4</sup>*Institut de Física d'Altes Energies, ICREA, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain*

<sup>5</sup>*Baylor University, Waco, Texas 76798, USA*

<sup>6a</sup>*Istituto Nazionale di Fisica Nucleare Bologna, I-40127 Bologna, Italy*

<sup>6b</sup>*University of Bologna, I-40127 Bologna, Italy*

<sup>7</sup>*University of California, Davis, Davis, California 95616, USA*

<sup>8</sup>*University of California, Irvine, Irvine, California 92627, USA*

<sup>9</sup>*University of California, Los Angeles, Los Angeles, California 90024, USA*

<sup>10</sup>*Instituto de Física de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*

<sup>11</sup>*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*

<sup>12</sup>*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA*

<sup>13</sup>*Comenius University, 842 48 Bratislava, Slovakia; Institute of Experimental Physics, 040 01 Kosice, Slovakia*

<sup>14</sup>*Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*

<sup>15</sup>*Duke University, Durham, North Carolina 27708, USA*

<sup>16</sup>*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*

<sup>17</sup>*University of Florida, Gainesville, Florida 32611, USA*

<sup>18</sup>*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*

<sup>19</sup>*University of Geneva, CH-1211 Geneva 4, Switzerland*

<sup>20</sup>*Glasgow University, Glasgow G12 8QQ, United Kingdom*

<sup>21</sup>*Harvard University, Cambridge, Massachusetts 02138, USA*

<sup>22</sup>*Division of High Energy Physics, Department of Physics, University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland*

<sup>23</sup>*University of Illinois, Urbana, Illinois 61801, USA*

<sup>24</sup>*The Johns Hopkins University, Baltimore, Maryland 21218, USA*

<sup>25</sup>*Institut für Experimentelle Kernphysik, Karlsruhe Institute of Technology, D-76131 Karlsruhe, Germany*

<sup>26</sup>*Center for High Energy Physics: Kyungpook National University, Daegu 702-701, Korea;*

*Seoul National University, Seoul 151-742, Korea; Sungkyunkwan University, Suwon 440-746, Korea;*

*Korea Institute of Science and Technology Information, Daejeon 305-806, Korea;*

*Chonnam National University, Gwangju 500-757, Korea;*

*Chonbuk National University, Jeonju 561-756, Korea*

<sup>27</sup>*Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*

<sup>28</sup>*University of Liverpool, Liverpool L69 7ZE, United Kingdom*

<sup>29</sup>*University College London, London WC1E 6BT, United Kingdom*

<sup>30</sup>*Centro de Investigaciones Energeticas Medioambientales y Tecnológicas, E-28040 Madrid, Spain*

<sup>31</sup>*Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

<sup>32</sup>*Institute of Particle Physics, McGill University, Montréal, Québec, Canada H3A 2T8;*

*Simon Fraser University, Burnaby, British Columbia, Canada V5A 1S6; University of Toronto, Toronto, Ontario, Canada M5S 1A7; and TRIUMF, Vancouver, British Columbia, Canada V6T 2A3*

<sup>33</sup>*University of Michigan, Ann Arbor, Michigan 48109, USA*

<sup>34</sup>*Michigan State University, East Lansing, Michigan 48824, USA*

<sup>35</sup>*Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia*

- <sup>36</sup>University of New Mexico, Albuquerque, New Mexico 87131, USA  
<sup>37</sup>The Ohio State University, Columbus, Ohio 43210, USA  
<sup>38</sup>Okayama University, Okayama 700-8530, Japan  
<sup>39</sup>Osaka City University, Osaka 588, Japan  
<sup>40</sup>University of Oxford, Oxford OX1 3RH, United Kingdom  
<sup>41a</sup>Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, I-35131 Padova, Italy  
<sup>41b</sup>University of Padova, I-35131 Padova, Italy  
<sup>42</sup>University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA  
<sup>43a</sup>Istituto Nazionale di Fisica Nucleare Pisa, I-56127 Pisa, Italy  
<sup>43b</sup>University of Pisa, I-56127 Pisa, Italy  
<sup>43c</sup>University of Siena, I-56127 Pisa, Italy  
<sup>43d</sup>Scuola Normale Superiore, I-56127 Pisa, Italy  
<sup>44</sup>University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA  
<sup>45</sup>Purdue University, West Lafayette, Indiana 47907, USA  
<sup>46</sup>University of Rochester, Rochester, New York 14627, USA  
<sup>47</sup>The Rockefeller University, New York, New York 10065, USA  
<sup>48a</sup>Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1, I-00185 Roma, Italy  
<sup>48b</sup>Sapienza Università di Roma, I-00185 Roma, Italy  
<sup>49</sup>Rutgers University, Piscataway, New Jersey 08855, USA  
<sup>50</sup>Texas A&M University, College Station, Texas 77843, USA  
<sup>51a</sup>Istituto Nazionale di Fisica Nucleare Trieste/Udine, I-34100 Trieste, I-33100 Udine, Italy  
<sup>51b</sup>University of Udine, I-33100 Udine, Italy  
<sup>52</sup>University of Tsukuba, Tsukuba, Ibaraki 305, Japan  
<sup>53</sup>Tufts University, Medford, Massachusetts 02155, USA  
<sup>54</sup>University of Virginia, Charlottesville, Virginia 22906, USA  
<sup>55</sup>Waseda University, Tokyo 169, Japan  
<sup>56</sup>Wayne State University, Detroit, Michigan 48201, USA  
<sup>57</sup>University of Wisconsin, Madison, Wisconsin 53706, USA  
<sup>58</sup>Yale University, New Haven, Connecticut 06520, USA  
(Received 5 December 2011; published 26 January 2012)

---

<sup>a</sup>Deceased

- <sup>b</sup>Visiting from Istituto Nazionale di Fisica Nucleare, Sezione di Cagliari, 09042 Monserrato (Cagliari), Italy  
<sup>c</sup>Visiting from University of California, Irvine, Irvine, CA 92697, USA  
<sup>d</sup>Visiting from University of California, Santa Barbara, Santa Barbara, CA 93106, USA  
<sup>e</sup>Visiting from University of California, Santa Cruz, Santa Cruz, CA 95064, USA  
<sup>f</sup>Visiting from Institute of Physics, Academy of Sciences of the Czech Republic, Czech Republic  
<sup>g</sup>Visiting from CERN, CH-1211 Geneva, Switzerland  
<sup>h</sup>Visiting from Cornell University, Ithaca, NY 14853, USA  
<sup>i</sup>Visiting from University of Cyprus, Nicosia CY-1678, Cyprus  
<sup>j</sup>Visiting from Office of Science, U.S. Department of Energy, Washington, DC 20585, USA  
<sup>k</sup>Visiting from University College Dublin, Dublin 4, Ireland  
<sup>l</sup>Visiting from ETH, 8092 Zurich, Switzerland  
<sup>m</sup>Visiting from University of Fukui, Fukui City, Fukui Prefecture, Japan 910-0017  
<sup>n</sup>Visiting from Universidad Iberoamericana, Mexico D.F., Mexico  
<sup>o</sup>Visiting from University of Iowa, Iowa City, IA 52242, USA  
<sup>p</sup>Visiting from Kinki University, Higashi-Osaka City, Japan 577-8502  
<sup>q</sup>Visiting from Kansas State University, Manhattan, KS 66506, USA  
<sup>r</sup>Visiting from Korea University, Seoul, 136-713, Korea  
<sup>s</sup>Visiting from University of Manchester, Manchester M13 9PL, United Kingdom  
<sup>t</sup>Visiting from Queen Mary, University of London, London, E1 4NS, United Kingdom  
<sup>u</sup>Visiting from University of Melbourne, Victoria 3010, Australia  
<sup>v</sup>Visiting from Muons, Inc., Batavia, IL 60510, USA  
<sup>w</sup>Visiting from Nagasaki Institute of Applied Science, Nagasaki, Japan  
<sup>x</sup>Visiting from National Research Nuclear University, Moscow, Russia  
<sup>y</sup>Visiting from Northwestern University, Evanston, IL 60208, USA  
<sup>z</sup>Visiting from University of Notre Dame, Notre Dame, IN 46556, USA  
<sup>aa</sup>Visiting from Universidad de Oviedo, E-33007 Oviedo, Spain  
<sup>bb</sup>Visiting from CNRS-IN2P3, Paris, F-75205 France, USA  
<sup>cc</sup>Visiting from Texas Tech University, Lubbock, TX 79609, USA  
<sup>dd</sup>Visiting from Universidad Tecnica Federico Santa Maria, 110v Valparaiso, Chile  
<sup>ee</sup>Visiting from Yarmouk University, Irbid 211-63, Jordan

We present a search for new phenomena in events with two reconstructed  $Z$  bosons and large missing transverse momentum, sensitive to processes  $p\bar{p} \rightarrow X_2 X_2 \rightarrow Z Z X_1 X_1$ , where  $X_2$  is an unstable particle decaying as  $X_2 \rightarrow Z X_1$  and  $X_1$  is undetected. The particles  $X_1$  and  $X_2$  may be, among other possibilities, fourth-generation neutrinos or supersymmetric particles. We study the final state in which one  $Z$  boson decays to two charged leptons and the second decays hadronically. In data corresponding to an integrated luminosity of  $4.2 \text{ fb}^{-1}$  from proton-antiproton collisions recorded by the CDF II detector at the Tevatron, with center-of-mass energy of 1.96 TeV, we find agreement between data and standard-model backgrounds. We calculate 95% confidence level upper limits on the cross section of the process  $p\bar{p} \rightarrow X_2 X_2 \rightarrow Z Z X_1 X_1$  ranging from 50 fb to 1 pb, depending on the masses of  $X_1$  and  $X_2$ .

DOI: 10.1103/PhysRevD.85.011104

PACS numbers: 12.60.-i, 13.85.Rm, 14.65.-q, 14.80.-j

A natural extension to the standard model of particle physics is a fourth generation of quarks and leptons. The inclusion of a fourth generation provides a source of  $CP$  violation in  $B_s$  decays and can accommodate a heavy Higgs boson [1,2]. Searches for fourth-generation quarks at the Fermilab Tevatron have constrained the mass of up-type quarks ( $u_4$ ), that decay as  $u_4 \rightarrow Wq$ , where  $q$  is a generic down-type quark, to be  $m_{u_4} > 340 \text{ GeV}/c^2$  at 95% confidence level (CL) [3], while limits on the mass of down-type quarks ( $d_4$ ) decaying via  $d_4 \rightarrow Wt$  are  $m_{d_4} > 372 \text{ GeV}/c^2$  at 95% CL [4].

Following the trend of mass hierarchy in the standard model, the least massive and therefore most accessible particle of this fourth generation may be the neutrino. Such a neutrino need not be solely a Dirac or Majorana state, but may be a mixture of the two [5]. This leads to two mass eigenstates  $N_1$  and  $N_2$ , where  $N_2$  is the unstable heavy eigenstate and  $N_1$  is the stable and least massive eigenstate of the fourth-generation neutrinos. These particles would partially evade the neutrino mass constraints from  $Z$  width studies at LEP [6].

The dominant production mechanism of  $N_1$  would be via a Drell-Yan process,  $p\bar{p} \rightarrow Z/\gamma^* \rightarrow N_2 N_2 \rightarrow N_1 Z N_1 Z$ , giving a final state of two  $Z$  bosons and large missing transverse momentum. This signature is shared by several other interesting new physics processes, most notably supersymmetric production,  $\chi_2^0 \chi_2^0 \rightarrow Z \chi_1^0 Z \chi_1^0$ , where  $\chi_1^0$  and  $\chi_2^0$  are neutralinos. We consider the mode in which one  $Z$  decays hadronically and the other decays leptonically, giving a detector signature of two charged leptons, two jets and large missing transverse momentum. For this search we use  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96 \text{ TeV}$  corresponding to  $4.2 \text{ fb}^{-1}$  of integrated luminosity collected by the CDF II detector.

Events were recorded by CDF II [7,8], a general-purpose detector designed to study collisions at the Fermilab Tevatron  $p\bar{p}$  collider. The CDF II detector is composed of a charged-particle tracking system immersed in a 1.4 T magnetic field consisting of a silicon microstrip tracker and a drift chamber. Electromagnetic and hadronic calorimeters surround the tracking system and measure particle energies. Drift chambers located outside the calorimeters detect muons.

The data acquisition system is triggered by  $e$  or  $\mu$  candidates with transverse momentum  $p_T$ , greater than  $18 \text{ GeV}/c$ . We retain electron and muon candidates with pseudorapidity [8]  $|\eta| < 1.1$ ,  $p_T \geq 20 \text{ GeV}/c$  and that satisfy the standard CDF identification requirements [9]. For muons, the track fit  $\chi^2$  per degree of freedom is used to reject poorly fit tracks likely resulting from charged pion and kaon decays in flight. Electrons from photon conversions are suppressed by rejecting electron candidates with a nearly collinear intersecting reconstructed track. Jets are reconstructed in the calorimeter using the JETCLU [10] algorithm with a clustering radius of 0.4 in azimuth-pseudorapidity space. Measured jet energies are corrected to account for  $\eta$ -dependent variations in detector response, calorimeter coverage, and the expected contribution from additional  $p\bar{p}$  interactions in the same event [11]. Jets are selected if they have  $p_T \geq 15 \text{ GeV}/c$  and  $|\eta| < 2.4$ . Missing transverse energy [12],  $\cancel{E}_T$ , is reconstructed using calorimeter and muon information including the corrections described above.

To isolate the  $ZZ$  signature, we require two opposite-charge, same-flavor lepton candidates ( $e$  or  $\mu$ ) with  $p_T > 20 \text{ GeV}/c$  for which the lepton-pair invariant mass is consistent with decay from a  $Z$  boson:  $m_{\ell\ell} [76, 106] \text{ GeV}/c^2$ . Additionally, we require at least two jets, each with  $p_T > 15 \text{ GeV}/c$  and  $|\eta| < 2.4$ , and without identified secondary vertices resulting from  $b$ -hadron decay [13]. The  $ZZ + \cancel{E}_T$  signature has the further requirement of large  $\cancel{E}_T$ , varying with hypothetical  $N_1$  and  $N_2$  masses, as shown in Table II.

The dominant background in the resulting sample is production of a  $Z$  boson in association with two jets from initial state radiation. We model this background using ALPGEN [16] to describe the hard process and PYTHIA [17] for the showering and hadronization. This background is strongly suppressed in events with large missing transverse momentum, as shown in Fig. 1 and Table I, and is distinguished from the signal by the lack of a resonance in the dijet mass,  $m_{jj}$ .

The second largest expected background is due to  $W$  boson production in association with three jets from initial state radiation, where one jet is wrongly reconstructed as a lepton. We model this using an independent sample of






FIG. 1. Distribution of missing transverse momentum in events with the ZZ signature, for expected backgrounds and observed data.

events containing jets likely to mimic leptons, following Ref. [18]. Additional backgrounds result from standard-model production of two gauge bosons, including ZZ, WW, and WZ, as well as  $t\bar{t} \rightarrow WbWb$ , which are all modeled using PYTHIA.

To isolate the double-resonance nature of the  $ZZ + \cancel{E}_T$  signature, we calculate the distance from the Z boson reconstructed mass in the  $m_{\ell\ell} - m_{jj}$  mass plane, accounting for the relative difference in the resolutions between the leptons and jets as well as the observed bias in reconstructed  $m_{jj}$ , using the variable

$$\Delta m = \sqrt{\left(\frac{m_{\ell\ell} - m_{Z \rightarrow \ell\ell}}{g_{\ell\ell}}\right)^2 + \left(\frac{m_{jj} - m_{Z \rightarrow jj}}{g_{jj}}\right)^2}, \quad (1)$$

where  $m_{\ell\ell}(m_{jj})$  is the reconstructed lepton (jet) pair mass, compared to the reference  $m_{Z \rightarrow \ell\ell} = 91.6 \text{ GeV}/c^2$

TABLE II. Acceptance of the  $ZZ + \cancel{E}_T$  selection for varying thresholds in  $\cancel{E}_T$  optimized for each point in the  $M_{N_1}, M_{N_2}$  mass plane. Also shown are the median expected and observed 95% CL upper limits on the cross section ( $\sigma_{N_2}$ ) in data with  $4.2 \text{ fb}^{-1}$  of integrated luminosity, as well as the theoretical prediction [14,15].

$M_{N_1}, M_{N_2}$ [GeV/ $c^2$ ]	$\cancel{E}_T$ cut [GeV]	Acceptance [%]	$\sigma_{N_2}$ [fb] Theory	$\sigma_{N_2}$ [fb] Exp./Obs. Limit
75, 175	37	0.99	0.51	511/702
75, 200	68	1.02	0.21	292/369
125, 225	36	0.85	0.16	684/1088
75, 225	92	0.93	0.081	156/273
75, 275	118	1.01	0.015	94/132
125, 300	119	1.06	0.013	99/138
175, 300	80	0.96	0.022	171/315
125, 350	156	1.05	0.003	75/48
225, 350	80	1.05	0.006	190/297
75, 350	167	1.06	0.001	71/55




FIG. 2. Distribution of the variable  $\Delta m$ , defined in the text, for expected background, observed data, and an example signal (scaled by  $10^4$ ) in data with  $4.2 \text{ fb}^{-1}$  of integrated luminosity. This example uses a missing transverse momentum threshold of  $\cancel{E}_T > 36 \text{ GeV}$ , optimized for this  $(M_{N_1}, M_{N_2})$  mass point; see Table II. Background uncertainties are statistical and systematic added in quadrature.

( $m_{Z \rightarrow jj} = 85.3 \text{ GeV}/c^2$ ) found in simulated events. To account for the superior lepton resolution, the dilepton and dijet mass differences are scaled by factors related to the resolutions:  $g_{\ell\ell} = 10 \text{ GeV}/c^2$ ,  $g_{jj} = 15 \text{ GeV}/c^2$ . The uncertainties of these reference values are small, and may be neglected. The distribution of  $\Delta m$  for data and simulated background and signal is shown in Fig. 2.

We model the production of the  $N_2$  signal and its subsequent decay into  $N_1$  over a grid of masses in the  $(M_{N_1}, M_{N_2})$  plane using MADGRAPH [14] with the CTEQ5L [19] parton distribution functions; PYTHIA [17] is used for the showering and hadronization. To suppress the large backgrounds expected from standard-model sources, we require large  $\cancel{E}_T$ ; as the expected magnitude of missing transverse momentum depends strongly on  $M_{N_1}$  and  $M_{N_2}$ , we vary the selection threshold of  $\cancel{E}_T$  to optimize for sensitivity at each  $(M_{N_1}, M_{N_2})$  pair considered, as seen in Table II. The acceptance for each mass point can be seen in Fig. 3. For each point in the mass grid, we form template histograms as a function of  $\Delta m$  for the expected signal and background, as displayed in Fig. 2.

In addition to the templates formed for the nominal expectation, we form alternate templates that incorporate the effects of systematic uncertainties under  $\pm 1\sigma$  variation. Fitting to these templates using the maximum likelihood method, we extract the best-fit signal cross section,  $\sigma_{N_2}$ . Systematic uncertainties affecting the shapes of templates, including uncertainty in the jet energy scale [11], QCD radiation, parton distribution functions,  $Q^2$  (square of momentum transfer in the interaction) and uncertainty in lepton energy resolution, are accounted for as nuisance parameters in our likelihood. The dominant source of systematic uncertainty in this analysis is uncertainty in the jet energy scale (40%), which can significantly modify




FIG. 3. Acceptance of the  $ZZ + \cancel{E}_T$  signature, including  $BR(ZZ \rightarrow \ell\ell qq)$ , as a function of the masses of the fourth-generation neutrinos,  $N_1$  and  $N_2$ . The threshold in  $\cancel{E}_T$  is optimized at each point on a grid in this plane. Linear interpolation is performed between the grid points. The apparent structure in the plot results from statistical fluctuation.

the number of jets in background processes that pass the  $p_T$  threshold, the location of the  $m_{jj}$  resonance in the signal process, and the measured  $\cancel{E}_T$  in an event. The second largest systematic uncertainty is due to uncertainty on the theoretical normalization of the background rates (10%). Finally, we apply the unified ordering principle [20] for the Neyman construction to create confidence intervals in the true value of  $\sigma_{N_2}$  for each  $N_2, N_1$  mass point.

We find the candidate events in the data to be consistent with expected standard-model backgrounds and thus set upper limits at 95% CL on the cross section for  $p\bar{p} \rightarrow N_2 N_2 \rightarrow N_1 Z N_1 Z$ . Theoretical cross sections for each mass point are presented in Table II, along with their respective expected and observed limits in our data sample. The expected and observed cross section limits can be seen in Fig. 4 and Table II.

In summary, we have performed the first search for new phenomena in events with two reconstructed  $Z$  bosons and

TABLE I. Expected number of events for each source of background to the  $ZZ \rightarrow \ell^+ \ell^- jj$  and  $ZZ + X_1 X_1 \rightarrow \ell^+ \ell^- jj + \cancel{E}_T$  signatures, as well as the observed event yield in data with  $4.2 \text{ fb}^{-1}$  of integrated luminosity. The threshold in  $\cancel{E}_T$  is optimized as a function of the  $N_1, N_2$  masses; one example ( $N_1 = 125 \text{ GeV}/c^2$ ,  $N_2 = 225 \text{ GeV}/c^2$ ) is shown here. Uncertainties shown include both systematic and statistical uncertainty added in quadrature.

Process	$\ell^+ \ell^- jj$	$\ell^+ \ell^- jj$ and $\cancel{E}_T > 36 \text{ GeV}$
$WW$	$4.4 \pm 1.3$	$2.7 \pm 0.8$
$t\bar{t}$	$14.8 \pm 3.0$	$11.6 \pm 2.3$
$W + \text{jets}$	$36.1 \pm 16.7$	$21.7 \pm 12.6$
$ZZ$	$99.4 \pm 20.5$	$4.2 \pm 0.9$
$WZ$	$105.6 \pm 22.1$	$5.2 \pm 1.1$
$Z + \text{jets}$	$10\,171 \pm 4422$	$94.6 \pm 38.5$
Total	$10\,432 \pm 4485$	$140.0 \pm 40.6$
Data	10 199	152




FIG. 4. Upper limit at 95% CL on the cross section of  $p\bar{p} \rightarrow N_2 N_2 \rightarrow N_1 Z N_1 Z$  in data with  $4.2 \text{ fb}^{-1}$  of integrated luminosity as a function of the masses of  $N_1$  and  $N_2$ . Top panel shows median expected limits; bottom panel shows observed limits; see Table II.

large missing transverse momentum. This signature is sensitive to processes  $p\bar{p} \rightarrow X_2 X_2 \rightarrow ZZ X_1 X_1$ , where  $X_2$  is an unstable particle decaying as  $X_2 \rightarrow ZX_1$  and  $X_1$  being undetected. The particles  $X_1$  and  $X_2$  may be, among other possibilities, fourth-generation neutrinos or supersymmetric particles. A specific model in which  $X_2$  and  $X_1$  are fourth-generation neutrinos is used without loss of generality. In the final state in which one  $Z$  boson decays to two charged leptons and the second decays hadronically, we find agreement between the data and the standard-model expectation using data from proton-antiproton collisions with  $4.2 \text{ fb}^{-1}$  of integrated luminosity. Based on the results in Table II, we report 95% CL upper limits on the cross section of the process  $p\bar{p} \rightarrow X_2 X_2 \rightarrow ZZ X_1 X_1$  ranging from 50 fb to 1 pb depending on the masses of  $X_1$  and  $X_2$ .

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the

Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the Korean World Class University Program, the National Research Foundation of Korea; the Science and Technology Facilities Council and the Royal Society,

UK; the Institut National de Physique Nucleaire et Physique des Particules/CNRS; the Russian Foundation for Basic Research; the Ministerio de Ciencia e Innovación, and Programa Consolider-Ingenio 2010, Spain; the Slovak R&D Agency; and the Academy of Finland.

- 
- [1] B. Holdom, W.S. Hou, T. Hurth, M.L. Mangano, S. Sultansoy, and G. Unel, *PMC Phys. A* **3**, 4 (2009).
  - [2] A.K. Alok, A. Dighe, and D. London, *Phys. Rev. D* **83**, 073008 (2011).
  - [3] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **107**, 261801 (2011).
  - [4] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **106**, 141803 (2011).
  - [5] L. Carpenter, [arXiv:1010.5502v1](#).
  - [6] P. Achard *et al.* (L3 Collaboration), *Phys. Lett. B* **517**, 75 (2001).
  - [7] D. Acosta *et al.* (CDF Collaboration), *Phys. Rev. D* **71**, 032001 (2005).
  - [8] CDF uses a cylindrical coordinate system with the  $z$  axis along the proton beam axis. Pseudorapidity is  $\eta \equiv -\ln(\tan(\theta/2))$ , where  $\theta$  is the polar angle relative to the proton beam direction, and  $\phi$  is the azimuthal angle while  $p_T = |p| \sin\theta$ ,  $E_T = E \sin\theta$ .
  - [9] A. Abulencia *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **97**, 082004 (2006); D. Acosta *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **94**, 091803 (2005).
  - [10] F. Abe *et al.* (CDF Collaboration), *Phys. Rev. D* **45**, 1448 (1992).
  - [11] A. Bhatti *et al.*, *Nucl. Instrum. Methods* **566**, 375 (2006).
  - [12] Missing transverse momentum,  $\cancel{E}_T$ , is defined as the vector  $-\sum_i E_T^i \vec{n}_i$ , where  $E_T^i$  are the magnitudes of transverse momentum contained in each calorimeter tower  $i$ , and  $\vec{n}_i$  is the unit vector from the interaction vertex to the tower in the transverse ( $x, y$ ) plane.
  - [13] D. Acosta *et al.* (CDF Collaboration), *Phys. Rev. D* **71**, 052003 (2005).
  - [14] J. Alwall, M. Herquet, F. Maltoni, O. Mattelaer, and T. Stelzer, *J. High Energy Phys.* **06** (2011) 128.
  - [15] L. Carpenter (private communication).
  - [16] M. Mangano *et al.*, *J. High Energy Phys.* **01** (2007) 013.
  - [17] T. Sjostrand, S. Mrenna, P. Skands, *J. High Energy Phys.* **05** (2006) 026.
  - [18] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **105**, 251802 (2010).
  - [19] J. Pumplin *et al.* (CTEQ Collaboration), *J. High Energy Phys.* **07** (2002) 012.
  - [20] G.J. Feldman and R.D. Cousins, *Phys. Rev. D* **57**, 3873 (1998).